



National Aeronautics and Space  
Administration Goddard Earth Science Data  
Information and Services Center (GES DISC)

# README Document for the Carbon Monitoring System (CMS) Carbon Flux Data Sets

---

## Prepared By:

Thomas Hearty

---

GES DISC  
GSFC Code 610.2

---

Kevin Bowman

---

Jet Propulsion Laboratory, California  
Institute of Technology

---

## Reviewed By:

Elliot Sherman

---

GES DISC  
GSFC Code 610.2

---

## Revision History

<i>Revision Date</i>	<i>Changes</i>	<i>Author</i>
9/8/2017	Original Document	Thomas Hearty
10/5/2017	Added dataset description and references	Kevin Bowman

Goddard Earth Sciences Data and Information Services Center (GES DISC)  
<http://disc.gsfc.nasa.gov>  
NASA Goddard Space Flight Center  
Code 610.2  
Greenbelt, MD 20771 USA

Part of the research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004)

# Table of Contents

Revision History .....	2
1.0 Introduction.....	4
1.1 Description of the Data Sets.....	5
1.2 Carbon Monitoring System (CMS) Description .....	5
1.3 Data Disclaimer and Digital Object Identifier (DOI).....	6
2.0 Data Organization .....	6
2.1 File Naming Convention .....	6
2.2 File Format and Structure.....	7
3.0 Data Contents.....	7
4.0 Options for Reading the Data .....	12
4.1 Programming Languages.....	12
4.2 Command Line Utility .....	12
4.3 A tool for simple visualization .....	12
5.0 Data Services .....	13
6.0 Acknowledgments.....	13
7.0 References.....	14

# 1.0 Introduction

This document provides basic information for using the 8 Carbon Monitoring System (CMS) datasets listed in Table 1.

**Table 1.** Datasets in this collection.

Dataset Title	Short Name/DOI
Carbon Monitoring System Flux for Fire L4 V1	CMSFluxFire 10.5067/3C1Y3EJB1E7Q
Carbon Monitoring System Flux for Posterior Fire Carbon L4 V1	CMSFluxFirepost 10.5067/N3HM4V0JZVLB
Carbon Monitoring System Flux for Fossil Fuel L4 V1	CMSFluxFossilfuel 10.5067/JC6BC3CPEJXQ
Carbon Monitoring System Flux for Shipping, Aviation, and Chemical Sources L4 V1	CMSFluxMISC 10.5067/RLT7JTCRJ11M
Carbon Monitoring System Flux from the Net Ecosystem Exchange L4 V1	CMSFluxNEE 10.5067/4ACY6GOWQ7BB
Carbon Monitoring System Flux for Ocean Carbon L4 V1	CMSFluxOcean 10.5067/96SSC2AOLE3Z
Carbon Monitoring System Flux for Posterior Total Carbon L4 V1	CMSFluxTotalpost 10.5067/QCBSYYY4CENP
Carbon Monitoring System Flux for Prior Total Carbon L4 V1	CMSFluxTotalprior 10.5067/F0JBNZ5QYWY6

## 1.1 Description of the Data Sets

These datasets contain global estimates of various components of the carbon cycle constrained by satellite observations through the Carbon Monitoring System Flux (CMS-Flux) carbon cycle data assimilation system as shown in Figure 1. A description of the methodology and technical details of the system can be found in Liu et al. 2014 and Bowman et al. 2017. The total CO<sub>2</sub> flux, which is the net sum of all carbon fluxes, from 2010-2016 are constrained with GOSAT v3.7b whereas total fluxes from 2015-2016 are constrained by OCO-2 v9r.




Figure: Carbon Monitoring System Flux (CMS-Flux) Framework. Satellite observations of surface data are integrated into a suite of anthropogenic (FFDAS), ocean (ECCO2-Darwin), and terrestrial (CASA-GFED) carbon cycle models. These are in turn used to compute surface fluxes that drive a chemistry and transport model (GEOS-Chem). Atmospheric observations of CO<sub>2</sub>, CO, CH<sub>4</sub> are ingested into an inverse model that computes posterior estimates of carbon surface fluxes. The combination of fluxes is used to attribute carbon and then reconcile those differences with prior carbon cycle models (from Bowman et al, 2017).

## 1.2 Carbon Monitoring System (CMS) Description

The NASA Carbon Monitoring System (CMS) is designed to make significant contributions in characterizing, quantifying, understanding, and predicting the evolution of global carbon sources and sinks through improved monitoring of carbon stocks and fluxes. The System will use the full range of NASA satellite observations and modeling/analysis capabilities to establish the accuracy, quantitative uncertainties, and utility of products for supporting national and international policy, regulatory, and management activities. CMS will maintain a global emphasis while providing finer scale regional information, utilizing space-based and surface-based data and will rapidly initiate generation and distribution of products both for user evaluation and to inform near-term policy development and planning.

## 1.3 Data Disclaimer and Digital Object Identifier (DOI)

The data sets may be acknowledged in publications using the Digital Object Identifiers listed in Table 1.

# 2.0 Data Organization

The data in all the files are organized on an equal angle grid in longitude and latitude but they have different spatial resolutions. The files also each contain one year of data.

## 2.1 File Naming Convention

The filenames are Described in Table 2 where YYYY should be replaced by the year.

**Table 2.** CMS Flux File Naming Conventions

Short Name	Filename pattern

CMS_Flux_Fire	CMS_Flux_Fire_YYYY_v2.nc4
CMS_Flux_FossilFuelPrior	CMS_Flux_FossilFuelPrior_YYYY_v2.nc4
CMSFluxMISC	CMSFluxMISC_v1.nc4
CMS_Flux_NBE	CMS_Flux_NBE_YYYY_v2.nc4
CMS_Flux_OceanPrior	CMS_Flux_OceanPrior_YYYY_v2.nc4
CMS_Flux_Total	CMS_Flux_Total_YYYY_v2.nc4

## 2.2 File Format and Structure

The files are stored in NetCDF-4 format.

## 3.0 Data Contents

The dimensions and variables of each of the CMS-Flux products are listed below.

CMS\_Flux\_Fire

dimensions:

```
lon = 72 ;
lat = 46 ;
time = 12 ;
```

variables:

```
float lon(lon) ;
    string lon:long_name = "longitude" ;
    string lon:units = "degrees_east" ;
float lat(lat) ;
    string lat:long_name = "latitude" ;
    string lat:units = "degrees_north" ;
float time(time) ;
    string time:long_name = "time" ;
    string time:units = "months since 2010-01-01" ;
float flux (time, lat, lon) ;
    string flux:long_name = " Posterior Biomass Burning Carbon Emissions (C)" ;
    string flux:units = "kg/km^2/s" ;
    long missing_value = -999;
float fluxunc (time, lat, lon) ;
    string fluxunc:long_name = " Posterior Biomass Burning Carbon Emissions
Uncertainty (C)" ;
    string fluxunc:units = "kg/km^2/s" ;
    long missing_value = -999;
float seconds_in_month (time);
    float :units = "seconds";
    string long_name = "Seconds in one month of a non-leap year.";
float area(lat) ;
    string area:long_name = "surface area per cell" ;
    string area:units = "km^2" ;
```

## CMS\_Flux\_FossilFuelPrior

dimensions:

```
lon = 72 ;  
lat = 46 ;  
time = 12;
```

variables:

```
float lon(lon) ;  
    string lon:long_name = "longitude" ;  
    string lon:units = "degrees_east" ;  
float lat(lat) ;  
    string lat:long_name = "latitude" ;  
    string lat:units = "degrees_north" ;  
float time(time) ;  
    string time:long_name = "time" ;  
    string time:units = "hours since 2010-01-01 00:00:0.0" ;  
float flux(time, lat, lon) ;  
    string flux:long_name = "Prior Fossil Fuel Carbon Emissions (C)" ;  
    string flux:units = "kg/km^2/s" ;  
    long missing_value = -999;  
float seconds_in_month (time);  
    float :units = "seconds";  
    string long_name = "Seconds in one month of a non-leap year.";  
float area(lat) ;  
    string Area:long_name = "surface area per cell" ;  
    string Area:units = "km^2" ;  
float seconds_in_month (time);  
    float :units = "seconds";
```

## CMSFluxMISC

dimensions:

```
lon = 72 ;  
lat = 46 ;  
lev = 47 ;  
time = 12 ;
```

variables:

```
float lon(lon) ;  
    string lon:long_name = "longitude" ;  
    string lon:units = "degrees_east" ;  
float lat(lat) ;  
    string lat:long_name = "latitude" ;
```

```

        string lat:units = "degrees_north" ;
float lev(lev) ;
        string lev:long_name = "GEOS-Chem vertical Level (1: surface)" ;
        string lev:units = "none" ;
float time(time) ;
        string time:long_name = "time" ;
        string time:units = "months since 0-01-01" ;
float Shipping(time, lat, lon) ;
        string Shipping:long_name = "Shipping Carbon Emission (ICOADS)" ;
        string Shipping:units = "kg/km^2/s" ;
float Area(lat) ;
        string Area:long_name = "surface area per cell" ;
        string Area:units = "km^2" ;
float Aviation(time, lev, lat, lon) ;
        string Aviation:long_name = "Aviation Carbon Emission (GEOSChem)" ;
        string Aviation:units = "kg/km^2/s" ;
float ChemicalSources(time, lev, lat, lon) ;
        string ChemicalSources:long_name = "Chemical Source Carbon Emission
(GEOSChem)" ;
        string ChemicalSources:units = "kg/km^2/s" ;
float seconds_in_month (time);
        float :units = "seconds";

```

#### CMS\_Flux\_NBE

dimensions:

```

lon = 72 ;
lat = 46 ;
time = 12 ;

```

variables:

```

float lon(lon) ;
        string lon:long_name = "longitude" ;
        string lon:units = "degrees_east" ;
float lat(lat) ;
        string lat:long_name = "latitude" ;
        string lat:units = "degrees_north" ;
float time(time) ;
        string time:long_name = "time" ;
        string time:units = "hours since 2010-01-01 00:00:0.0" ;
float flux(time, lat, lon) ;
        string flux:long_name = "Posterior Net Biome Carbon Exchange (C)" ;
        string flux:units = "kg/km^2/s" ;
        long missing_value = -999;
float fluxunc(time, lat, lon) ;

```

```

(C)";
    string flux:long_name = "Posterior Net Biome Carbon Exchange Uncertainty
    string flux:units = "kg/km^2/s" ;
    long missing_value = -999;
float area(lat) ;
    string area:long_name = "surface area per cell" ;
    string area:units = "km^2" ;
float seconds_in_month (time);
    float :units = "seconds";

```

#### CMS\_Flux\_NBEPrior

dimensions:

```

lon = 72 ;
lat = 46 ;
time = 12 ;

```

variables:

```

float lon(lon) ;
    string lon:long_name = "longitude" ;
    string lon:units = "degrees_east" ;
float lat(lat) ;
    string lat:long_name = "latitude" ;
    string lat:units = "degrees_north" ;
float time(time) ;
    string time:long_name = "time" ;
    string time:units = "hours since 2010-01-01 00:00:0.0" ;
float flux(time, lat, lon) ;
    string flux:long_name = "Prior Net Biome Carbon Exchange (C)";
    string flux:units = "kg/km^2/s" ;
    long missing_value = -999;
float Area(lat) ;
    string Area:long_name = "surface area per cell" ;
    string Area:units = "km^2" ;
float seconds_in_month (time);
    float :units = "seconds";

```

#### CMS\_Flux\_OceanPrior

dimensions:

```

lon = 72 ;
lat = 46 ;
time = 12 ;

```

variables:



```

float lon(lon) ;
    string lon:long_name = "longitude" ;
    string lon:units = "degrees_east" ;
float lat(lat) ;
    string lat:long_name = "latitude" ;
    string lat:units = "degrees_north" ;
float time(time) ;
    string time:long_name = "time" ;
    string time:units = "hours since 2010-01-01 00:00:0.0" ;
float flux(time, lat, lon) ;
    string Ocean:long_name = "Prior Ocean-Atmosphere Carbon Flux" ;
    string Ocean:units = "kg/km^2/s" ;
    long missing_value = -999
float area(lat) ;
    string Area:long_name = "surface area per cell" ;
    string Area:units = "km^2" ;
float seconds_in_month (time);
    float :units = "seconds";

```

#### CMS\_Flux\_Total

dimensions:

```

lon = 72 ;
lat = 46 ;
time = 12 ;

```

variables:

```

float lon(lon) ;
    string lon:long_name = "longitude" ;
    string lon:units = "degrees_east" ;
float lat(lat) ;
    string lat:long_name = "latitude" ;
    string lat:units = "degrees_north" ;
float time(time) ;
    string time:long_name = "time" ;
    string time:units = "months since 2010-01-01" ;
float flux(time, lat, lon) ;
    string flux:long_name = "Posterior Total Surface-Atmosphere Exchange Carbon
(C)" ;
    string flux:units = "kg/km^2/s" ;
    long missing_value = -999
float area(lat) ;
    string area:long_name = "surface area per cell" ;
    string area:units = "km^2" ;
float seconds_in_month (time);
    float :units = "seconds";

```

## 4.0 Options for Reading the Data

### 4.1 Programming Languages

The data can be read using major programming languages such as Fortran, C, Java, IDL, Matlab, and Python.

### 4.2 Command Line Utility

#### **ncdump**

The ncdump tool can be used as a simple browser for NetCDF and HDF data files, to display the dimension names and sizes; variable names, types, and shapes; attribute names and values; and optionally, the values of data for all variables or selected variables in a netCDF file. The most common use of ncdump is with the `-h` option, in which only the header information is displayed.

```
ncdump [-c|-h] [-v ...] [[-b|-f] [c|f]] [-l len] [-n name] [-d n[,n]] filename
```

Options/Arguments:

`[-c]` Coordinate variable data and header information

`[-h]` Header information only, no data

`[-v var1[,...]]` Data for variable(s) `<var1>`,... only data

`[-f [c|f]]` Full annotations for C or Fortran indices in data

`[-l len]` Line length maximum in data section (default 80)

`[-n name]` Name for netCDF (default derived from file name)

`[-d n[,n]]` Approximate floating-point values with less precision filename File name of input netCDF file

### 4.3 A tool for simple visualization

**Panoply**, developed at the Goddard Institute for Space Studies (GISS), is compliant with NetCDF Climate and Forecast (CF) Metadata Convention that is gaining popularity. A strength of the tool is that data can be previewed “remotely” over the network – i.e. user can preview file content of HDF files stored on a remote site, without downloading them. Panoply is available from GISS:

<http://www.giss.nasa.gov/tools/panoply/>

## 5.0 Data Services

Data services and access methods can be found on the dataset landing page for each product:

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxFire\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxFire_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxFirepost\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxFirepost_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxFossilfuel\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxFossilfuel_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxMISC\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxMISC_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxNEE\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxNEE_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxOcean\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxOcean_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxTotalpost\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxTotalpost_1.html)

[http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxTotalprior\\_1.html](http://disc.sci.gsfc.nasa.gov/datacollection/CMSFluxTotalprior_1.html)

If you need assistance or wish to report a problem:

**Email:** [gsfc-help-disc@lists.nasa.gov](mailto:gsfc-help-disc@lists.nasa.gov)

**Voice:** 301-614-5224

**Fax:** 301-614-5268

**Address:**

Goddard Earth Sciences Data and Information Services Center NASA Goddard Space Flight  
Center Code 610.2 Greenbelt, MD 20771 USA

## 6.0 Acknowledgments

The distribution of this data is funded by the NASA Carbon Monitoring System CMS program.

## 7.0 References

Liu, J., K. Bowman, M. Lee, D. Henze, N. Bousserez, H. Brix, G. J. Collatz, D. Menemenlis, L. Ott, S. Pawson, D. Jones, and R. Nassar, Carbon monitoring system flux estimation and attribution: impact of ACOS-GOSAT XCO<sub>2</sub> sampling on the inference of terrestrial biospheric sources and sinks, *Tellus B*, 66(0), doi:<http://dx.doi.org/10.3402/tellusb.v66.22486>, 2014.

Bowman, K. W., J. Liu, A. A. Bloom, N. C. Parazoo, M. Lee, Z. Jiang, D. Menemenlis, M. M. Gierach, G. J. Collatz, K. R. Gurney, and D. Wunch, Global and Brazilian carbon response to El Niño Modoki 2011-2010, *Earth and Space Science*, 4, doi:[10.1002/2016EA000204](https://doi.org/10.1002/2016EA000204), 2017.